

Table of Contents

02.06.01 - IDAHO DEPARTMENT OF AGRICULTURE RULES PURE SEED LAW

| | |
|--|----|
| 000. -- 009. (RESERVED). | 2 |
| 010. METHODS OF SAMPLING -- GENERAL PROCEDURE. | 2 |
| 011. -- 049. (RESERVED). | 3 |
| 050. GERMINATION STANDARDS -- TABLE. | 3 |
| 051. -- 099. (RESERVED). | 4 |
| 100. GERMINATION STANDARDS FOR FLOWER SEEDS. | 4 |
| 101. -- 149. (RESERVED). | 10 |
| 150. GERMINATION STANDARDS FOR SEED IN HERMETICALLY SEALED CONTAINERS. | 10 |
| 151. -- 199. (RESERVED). | 12 |
| 200. NOXIOUS WEEDS. | 12 |
| 201. -- 249. (RESERVED). | 14 |
| 250. LABEL REQUIREMENTS OF SEEDS FOR SPROUTING. | 14 |
| 251. -- 299. (RESERVED). | 15 |
| 300. VIABILITY BY TZ%. | 15 |
| 301. -- 399. (RESERVED). | 15 |
| 400. METHODS OF TESTING. | 15 |
| 401. -- 999. (RESERVED). | 15 |

**IDAPA 02
TITLE 06
Chapter 01**

02.06.01 - IDAHO DEPARTMENT OF AGRICULTURE RULES PURE SEED LAW

000. -- 009. (RESERVED).

010. METHODS OF SAMPLING -- GENERAL PROCEDURE.

01. Sample. In order to secure a representative sample, equal portions shall be taken from evenly distributed parts of the quantity of seed or screenings to be sampled. Access shall be had to all parts of that quantity. When more than one trierful of seed is drawn from a bag, different paths shall be followed. When more than one handful is taken from a bag, the handfuls shall be taken from well-separated points. (7-1-93)

02. Free Flowing Seed. For free-flowing seed in bags or bulk, a probe or trier shall be used. For small free-flowing seed in bags a probe or trier long enough to sample all portions of the bag shall be used. (7-1-93)

03. Non-Free Flowing Seed. Non-free-flowing seed, such as certain grass seed, uncleaned seed or screenings difficult to sample with a probe or trier, shall be sampled by thrusting the hand into the bulk and withdrawing representative portions. The hand is inserted in an open position and the fingers are held closely together while the hand is being inserted and the portion withdrawn. (7-1-93)

04. Composite Samples. Composite samples shall be obtained to determine the quality of a lot of seed (i.e., percentages of pure seed, other crop seed, weed seed, inert matter and germination). Individual bag samples may be obtained to determine if the lot of seed is uniform. (7-1-93)

a. To determine if there is an obvious lack of uniformity of seed from which a composite sample is being obtained, each portion shall be examined and the portions shall then be combined to form a composite sample or samples. (7-1-93)

b. If the lot is found not to be uniform when obtaining a composite sample to determine its quality then additional individual bag samples shall be taken for the purpose of testing for uniformity. (7-1-93)

c. Such individual bag samples may also be taken for the purpose of testing for uniformity even though a composite sample has not previously been obtained. The identity of each individual bag sample must be maintained. (7-1-93)

05. Bulk. Bulk seed or screenings shall be sampled by inserting a long probe or thrusting the hand into the bulk, as circumstances require, in at least seven uniformly distributed parts of the quantity being sampled. At least as many trierfuls or handfuls shall be taken as the minimum which would be required for the same quantity of seed or screenings in bags of a size customarily used for such seed or screenings. (7-1-93)

06. Bags. (7-1-93)

a. For lots of six (6) bags or less, each bag shall be sampled. A total of at least five (5) trierfuls shall be taken. (7-1-93)

b. For lots of more than six (6) bags, five (5) bags plus at least 10 percent (10%) of the number of bags in the lots shall be sampled. (Round off numbers with decimals to the nearest whole number, raising five tenths (.5) to the next whole number.) Regardless of the lot size it is not necessary that more than thirty (30) bags be sampled. (7-1-93)

c. Samples shall be drawn from unopened bags except under circumstances where the identity of the seed has been preserved. (7-1-93)

07. Packets. In sampling seed in packets, entire unopened packets shall be taken. (7-1-93)

08. Size of Sample. (7-1-93)

- a. For composite sample to test for quality, the following are minimum weights for samples of seed to be submitted for analysis, test or examination (7-1-93)
- i. Two (2) ounces (approximately fifty five (55) grams) of grass seed not otherwise mentioned, white or alsike clover, or seeds not larger than these. (7-1-93)
 - ii. Five (5) ounces (approximately one hundred fifty (150) grams) of red or crimson clover, alfalfa, lespedezas, ryegrasses, bromegrasses, millet, flax, rape, or seeds of similar size. (7-1-93)
 - iii. One pound of Sudangrass, sorghum, proso, hemp seed or seeds of similar size. (7-1-93)
 - iv. Two (2) pounds (approximately one thousand (1,000) grams) of cereals, vetches, or seeds of similar or larger size. (7-1-93)
- b. For individual bag samples to test for uniformity. (7-1-93)
- i. The size of any individual bag sample to determine uniformity in a lot of seed shall be not less than the quantities set out in the "Rules and Regulations, under the Federal Seed Act" (53 Statute 1275) (Subsection 201.46). (7-1-93)
 - ii. If the sample drawn is larger than desired, it shall be thoroughly mixed before it is divided to the desired size. (7-1-93)
09. Forwarding and Receipt of Official Samples. Before being forwarded for analysis test or examination, the containers of official samples shall be properly sealed and identified. The containers of official samples shall be initialed and dated and the sample weighed by the person who breaks the seals. (7-1-93)

011. -- 049. (RESERVED).

050. GERMINATION STANDARDS -- TABLE.

Germination Standards for Vegetable Seeds, which shall be construed to include hard seed.

| | Percent | | Percent |
|----------------------------|---------|------------------|---------|
| Artichoke | 60 | Eggplant | 60 |
| Asparagus | 70 | Endive | 70 |
| Asparagusbean | 75 | Kale | 75 |
| Bean, garden | 70 | Kale, Chinese | 75 |
| Bean, lima | 70 | Kohlrabi | 75 |
| Bean, runner | 75 | Leek | 60 |
| Beet | 65 | Lettuce | 80 |
| Broadbean | 75 | Muskmelon | 75 |
| Broccoli | 75 | Mustard, India | 75 |
| Brussels sprouts | 70 | Mustard, spinach | 75 |
| Burdock, great | 60 | Okra | 50 |
| Cabbage | 75 | Onion | 70 |
| Cabbage, tronchuda | 75 | Onion, Welsh | 70 |
| Cantaloupe (See muskmelon) | | Pak-choi | 75 |

| | Percent | | Percent |
|-----------------|---------|----------------------|---------|
| Cardoon | 60 | Parsley | 60 |
| Carrot | 55 | Parsnip | 60 |
| Cauliflower | 75 | Pea | 80 |
| Celeriac | 55 | Pepper | 55 |
| Celery | 55 | Pumpkin | 75 |
| Chard, Swiss | 65 | Radish | 75 |
| Chicory | 65 | Rhubarb | 60 |
| Chinese Cabbage | 75 | Rutabaga | 75 |
| Chives | 50 | Salsify | 75 |
| Citron | 65 | Sorrel | 65 |
| Collards | 80 | Soybean | 75 |
| Corn, sweet | 75 | Spinach | 60 |
| Cornsalad | 70 | Spinach, New Zealand | 40 |
| Cowpea | 75 | Squash | 75 |
| Cress, garden | 75 | Tomato | 75 |
| Cress, upland | 60 | Tomato, husk | 50 |
| Cress, water | 40 | Turnip | 80 |
| Cucumber | 80 | Watermelon | 70 |
| Dandelion | 60 | | |

(7-1-93)

051. -- 099. (RESERVED).

100. GERMINATION STANDARDS FOR FLOWER SEEDS.

The kinds of flower seeds listed below are those for which standard testing procedures have been prescribed by the Association of Official Seed Analysts (AOSA) and which are required by the labeling provisions of Section 22-415, Idaho Code. The percentage listed opposite each kind is the germination standard for that kind. For the kinds marked with an asterisk, the percentage is arrived at by totaling the percent germination and percent hard seed. (7-1-93)

01. Table 1.

(7-1-93)

| Kind: | Percent: |
|---|----------|
| Achillea (The Pearl) - Achillea ptarmica | 50 |
| African Daisy - Dimorphotheca aurantiaca | 55 |
| African Violet - Saintpaulia spp. | 30 |
| Ageratum - Ageratum mexicanum | 60 |
| Agrostemma (rose champion) - Agrostemma coronaria | 65 |

| Kind: | Percent: |
|---|----------|
| Alyssum - Alyssum campactum, A. maritimum, A. procumbens, A. saxatile | 60 |
| Amaranthus - Amaranthus spp. | 65 |
| Anagalis (pimpernel) - Anagalis arvensis, Anagalis coerulea, Anagalis grandiflora | 60 |
| Anemone - Anemone coronaria, A. pulsatilla | 55 |
| Angel's Trumpet - Datura arborea | 60 |
| Arabis - Arabis alpina | 60 |
| Arctotis (African lilac daisy) - Arctotis grandis | 45 |
| Armeria - Armeria formosa | 55 |
| Asparagus, fern - Asparagus plumosus | 50 |
| Asparagus, sprenger - Asparagus sprengeri | 55 |
| Aster, China - Callistephus chinensis; except Pompon, Powderpuff, and Princess types | 55 |
| Aster, China - Callistephus chinensis; Pompon, Powderpuff, and Princess types | 50 |
| Aubretia - Aubretia deltoides | 45 |
| Baby Smilax - Asparagus asparagoides | 25 |
| Balsam - Impatiens balsamina | 70 |
| Begonia - Begonia fibrous rooted | 60 |
| Begonia - Begonia tuberous rooted | 50 |
| Bells of Ireland - Molucella laevis | 60 |
| Brachycome (swan river daisy) - Brachycome iberidifolia | 60 |
| Browallia - Browallia elata and B. speciosa | 65 |
| Buphthalmum (sunwheel) - Buphthalmum salicifolium | 60 |
| Calceolaria - Calceolaria spp. | 60 |
| Calendula - Calendula officinalis | 65 |
| California Poppy - Eschscholtzia californica | 60 |
| Calliopsis - Coreopsis bicolor, C. drummondi, C. elegans | 65 |
| Campanula: Canterbury Bells - Campanula medium | 60 |
| Cup and Saucer Bellflower - Campanula medium calycanthema | 60 |
| Carpathian Bellflower - Campanula carpatica | 50 |
| Peach Bellflower - Campanula persicifolia | 50 |
| Candytuft, Annual - Iberis amara, I. umbellata | 65 |
| Candytuft, Perennial - Iberis gibraltarica, I. sempervirens | 55 |
| *Castor Bean - Ricinus communis | 60 |

| Kind: | Percent: |
|--|----------|
| Cathedral Bells - <i>Cobaea scandens</i> | 65 |
| Celosia - <i>Celosia argentea</i> | 65 |
| Centaurea: Basket Flower - <i>Centaurea americana</i> , Cornflower - <i>C. cyanus</i> , Dusty Miller - <i>C. candidissima</i> , Royal Centaurea - <i>C. imperialis</i> , Sweet Sultan - <i>C. moschata</i> , Velvet Centaurea - <i>C. gymnocarpa</i> | 60 |
| Cerastium (snow-in-summer) - <i>Cerastium biebersteini</i> and <i>C. tomentosum</i> | 65 |
| Chinese Forget-me-not - <i>Cynoglossum amabile</i> | 55 |
| Chrysanthemum, Annual - <i>Chrysanthemum carinatum</i> , <i>C. coronarium</i> , <i>C. segetum</i> | 40 |
| Cineraria - <i>Senecio cruentus</i> | 60 |
| Clarkia - <i>Clarkia elegans</i> | 65 |
| Cleome - <i>Cleome gigantea</i> | 65 |
| Coleus - <i>Coleus blumei</i> | 65 |
| Columbine - <i>Aquilegia</i> spp. | 50 |
| Coral Bells - <i>Heuchera sanguinea</i> | 55 |
| Coreopsis, Perennial - <i>Coreopsis lanceolata</i> | 40 |
| Corn, ornamental - <i>Zea mays</i> | 75 |
| Cosmos: Sensation, Mammoth and Crested types - <i>Cosmos bipinnatus</i> ; Klondyke type - <i>C. sulhureus</i> | 65 |
| Crossandra - (<i>Crossandra infundibuliformis</i>) | 50 |
| Dahlia - <i>Dahlia</i> spp. | 55 |
| Daylily - <i>Hemerocallis</i> spp. | 45 |
| Delphinium, Perennial: Belladonna and Bellamosum types; Cardinal Larkspur - <i>Delphinium cardinale</i> ; Chinensis types; Pacific Giant, Gold Medal and other hybrids of <i>D. elatum</i> | 55 |
| Dianthus: Carnation - <i>Dianthus caryophyllus</i> | 60 |
| China Pinks - <i>Dianthus chinensis</i> , <i>heddewigi</i> , <i>heddensis</i> | 70 |
| Grass Pinks - <i>Dianthus plumarius</i> | 60 |
| Maiden Pinks - <i>Dianthus deltoides</i> | 60 |
| Sweet William - <i>Dianthus barbatus</i> | 70 |
| Sweet Wivelsfield - <i>Dianthus allwoodi</i> | 60 |
| Didiscus - (blue lace flower) - <i>Didiscus coerulea</i> | 65 |
| Doronicum (leopard's bane) - <i>Doronicum caucasicum</i> | 60 |
| Dracaena - <i>Dracaena indivisa</i> | 55 |

| Kind: | Percent: |
|--|----------|
| Dragon Tree - <i>Dracaena draco</i> | 40 |
| English Daisy - <i>Bellis perennis</i> | 55 |
| Flax - Golden flax (<i>Linum flavum</i>); Flowering flax <i>L. grandiflorum</i> ; Perennial flax, <i>L. perenne</i> | 60 |
| Flowering Maple - <i>Abutilon</i> spp. | 35 |
| Foxglove - <i>Digitalis</i> spp. | 60 |
| Gaillardia, Annual - <i>Gaillardia pulchella</i> ; <i>G. picta</i> ; Perennial - <i>G. grandiflora</i> | 45 |
| Gerbera (transvaal daisy) - <i>Gerbera jamesoni</i> | 60 |
| Geum - <i>Geum</i> spp. | 55 |
| Gilia - <i>Gilia</i> spp. | 65 |
| Gloriosa daisy (rudbeckia) - <i>Echinacea purpurea</i> and <i>Rudbeckia hirta</i> | 60 |
| Gloxinia - (<i>Sinningia speciosa</i>) | 40 |
| Godetia - <i>Godetia amoena</i> , <i>G. grandiflora</i> | 65 |
| Gourds: Yellow Flowered - <i>Cucurbita pepo</i> ; White Flowered - <i>Lagenaria siceraria</i> ; Dishcloth - <i>Luffa cylindrica</i> | 70 |
| Gypsophila: Annual Baby's Breath - <i>Gypsophila elegans</i> ; Perennial Baby's Breath - <i>G. paniculata</i> , <i>G. pacifica</i> , <i>G. repens</i> | 70 |
| Helenium - <i>Helenium autumnale</i> | 40 |
| Helichrysum - <i>Helichrysum monstrosum</i> | 60 |
| Heliopsis - <i>Heliopsis scabra</i> | 55 |
| Heliotrope - <i>Heliotropium</i> spp. | 35 |
| Helipterum (Acroclinium) - <i>Helipterum roseum</i> | 60 |
| Hesperis (sweet rocket) - <i>Hesperis matronalis</i> | 65 |
| *Hollyhock - <i>Althea rosea</i> | 65 |
| Hunnemania (mexican tulip poppy) - <i>Hunnemania fumariaefolia</i> | 60 |
| *Hyacinth bean - <i>Dolichos lablab</i> | 70 |
| Impatiens - <i>Impatiens holstii</i> , <i>I. sultani</i> | 55 |
| *Ipomea - Cypress Vine - <i>Ipomea quamoclit</i> ; Moonflower - <i>I. noctiflora</i> ; Morning Glories, Cardinal Climber, Hearts and Honey Vine - <i>Ipomea</i> spp. | 75 |
| Jerusalem cross (maltese cross) - <i>Lychnis chalconica</i> | 70 |

| Kind: | Percent: |
|---|----------|
| Job's Tears - <i>Coix lacrymajobi</i> | 70 |
| Kochia - <i>Kochia childsi</i> | 55 |
| Larkspur, Annual - <i>Delphinium ajacis</i> | 60 |
| Lantana - <i>Lantana camara</i> , <i>L. hybrida</i> | 35 |
| Lilium (regal lily) - <i>Lilium regale</i> | 50 |
| Linaria - <i>Linaria</i> spp. | 65 |
| Lobelia, Annual - <i>Lobelia erinus</i> | 65 |
| Lunaria, Annual - <i>Lunaria annua</i> | 65 |
| *Lupine - <i>Lupinus</i> spp. | 65 |
| Marigold - <i>Tagetes</i> spp. | 65 |
| Marvel of Peru - <i>Mirabilis jalapa</i> | 60 |
| Matricaria (feverfew) - <i>Matricaria</i> spp. | 60 |
| Mignonette - <i>Reseda odorata</i> | 55 |
| Myosotis - <i>Myosotis alpestris</i> , <i>M. oblongata</i> , <i>M. palustris</i> | 50 |
| Nasturtium - <i>Tropaeolum</i> spp. | 60 |
| Nemesia - <i>Nemesia</i> spp. | 65 |
| Nemophila - <i>Nemophila insignis</i> | 70 |
| Nemophila, spotted - <i>Nemophila maculata</i> | 60 |
| Nicotiana - <i>Nicotiana affinis</i> , <i>N. sanderae</i> , <i>N. sylvestris</i> | 65 |
| Nierembergia - <i>Nierembergia</i> spp. | 55 |
| Nigella - <i>Nigella damascena</i> | 55 |
| Pansy - <i>Viola tricolor</i> | 60 |
| Penstemon - <i>Penstemon barbatus</i> , <i>P. grandiflorus</i> , <i>P. laevigatus</i> , <i>P. pubescens</i> | 60 |
| Petunia - <i>Petunia</i> spp. | 45 |
| Phacelia - <i>Phacelia campanularia</i> , <i>P. minor</i> , <i>P. tanacetifolia</i> | 65 |
| Phlox, Annual - <i>Phlox drummondii</i> all types and varieties | 55 |
| Physalis - <i>Physalis</i> spp. | 60 |
| Platycodon (balloon flower) - <i>Platycodon grandiflorum</i> | 60 |
| Plumbago, cape - <i>Plumbago capensis</i> | 50 |
| Ponytail - <i>Beaucarnea recurvata</i> | 40 |
| Poppy: Shirley Poppy - <i>Papaver rhoeas</i> ; Iceland Poppy - <i>P. nudicaule</i> ; Oriental Poppy - <i>P. orientale</i> ; Tulip Poppy - <i>P. glaucum</i> | 60 |
| Portulace - <i>Portulaca grandiflora</i> | 55 |

| Kind: | Percent: |
|--|----------|
| Primula (Primrose) - Primula spp. | 50 |
| Pyrethrum (painted daisy) - Pyrethrum coccineum | 60 |
| Salpiglossis - Salpiglossis gloxinaeflora, S. sinuata | 60 |
| Salvia - Scarlet Sage - Salvia splendens; Mealycup Sage (blue bedder) - Salvia farinacea | 50 |
| Saponaria - Saponaria ocymoides, S. vaccaria | 60 |
| Scabiosa, Annual - Scabiosa atropurpurea | 50 |
| Scabiosa, Perennial - Scabiosa caucasica | 40 |
| Schizanthus - Schizanthus spp. | 60 |
| *Sensitive plant (mimosa) - Mimosa pudica | 65 |
| Shasta Daisy - Chrysanthemum maximum, C. leucanthemum | 65 |
| Silk Oak - Grevillea robusta | 25 |
| Snapdragon - Antirrhinum spp. | 55 |
| Solanum - Solanum spp. | 60 |
| Statice - Statice sinuata, S. suworonii (flower heads) | 50 |
| Stocks: Common - Mathiola incana; Evening Scented - Mathiola bicornis | 65 |
| Sunflower - Helianthus spp. | 70 |
| Sunrose - Helianthemum spp. | 30 |
| *Sweet Pea, Annual and Perennial other than dwarf bush - Lathyrus odoratus, L. latifolius | 75 |
| *Sweet Pea, dwarf bush - Lathyrus odoratus | 65 |
| Tahoka daisy - Machaeranthera tanacetifolia | 60 |
| Thunbergia - Thunbergia alata | 60 |
| Torch Flower - Tithonia speciosa | 70 |
| Torenia (wishbone flower) - Torenia fournieri | 70 |
| Tritoma Kniphofia spp. | 65 |
| Verbena, Annual - Verbena hybrida | 35 |
| Vinca - Vinca rosea | 60 |
| Viola - Viola cornuta | 55 |
| Virginian stocks - Malcolmia maritima | 65 |
| Wallflower - Cheiranthus allioni | 65 |
| Yucca (Adam's needle) - Yucca filamentosa | 50 |
| Zinnia (except Linearis and Creeping) - Zinnia angustifolia, Z. elegans, Z. grandiflora, Z. gracillima, Z. haegeana, Z. multiflora, Z. pumila | 65 |

| Kind: | Percent: |
|--|----------|
| Zinnia, Linearis and Creeping - Zinnia linearis, Sanvitalia procumbens | 50 |
| All other kinds | 50 |

(7-1-93)

02. Below Standard. A mixture of kinds of flower seeds will be considered to be below standard if the germination of any kind or combination of kinds constituting 25 percent (25%) or more of the mixture by number is below standard for the kind or kinds involved. (7-1-93)

101. -- 149. (RESERVED).

150. GERMINATION STANDARDS FOR SEED IN HERMETICALLY SEALED CONTAINERS.

The period of validity of germination tests is extended to the following period for seed packaged in hermetically sealed containers under conditions and label requirements set forth in this section. (7-1-93)

01. Germination Tests. The germination test for agricultural and vegetable seed shall have been completed within the following period, exclusive of the calendar month in which the test was completed, immediately prior to shipment, delivery, transportation or sales: (7-1-93)

a. In the case of agricultural or vegetable seeds shipped, delivered, transported or sold to a dealer for resale, 18 months; (7-1-93)

b. In the case of agricultural or vegetable seeds for sale or sold at retail, thirty-six (36) months. (7-1-93)

02. Conditions of Packaging. The following standards, requirements, conditions must be met before seed is considered to be hermetically sealed under the provisions of this Act: (7-1-93)

a. The seed was packaged within nine months after harvest. (7-1-93)

b. The container used does not allow water vapor penetration through any wall, including the seals, greater than five hundredths (0.05) gram of water per twenty-four (24) hours per one hundred (100) square inches of surface at one hundred degrees F. (100F) with a relative humidity on one side of ninety percent (90%) and on the other of zero percent (0%). Water vapor penetration or WVP is measured by the standards of the U.S. Bureau of Standards as-gm. H₂O/24 hr./100 sq. in./100 F/ 90% RHV. 0% RH. (7-1-93)

03. Moisture. The seed in the container does not exceed the percentage of moisture, on a wet weight basis, as listed below: (7-1-93)

a. Table 1. (7-1-93)

| Agricultural Seeds | Percent |
|---------------------|---------|
| Beet, field | 7.5 |
| Beet, sugar | 7.5 |
| Bluegrass, Kentucky | 6.0 |
| Clover, crimson | 8.0 |
| Fescue, red | 8.0 |
| Ryegrass, annual | 8.0 |

| Agricultural Seeds | Percent |
|------------------------------|---------|
| Ryegrass, perennial | 8.0 |
| All other agricultural seeds | 6.0 |
| Mixtures of above | 8.0 |

b. Table 2.

(7-1-93)

| Vegetable Seeds | Percent |
|------------------|---------|
| Bean, garden | 7.0 |
| Bean, lima | 7.0 |
| Beet | 7.5 |
| Broccoli | 5.0 |
| Brussels sprouts | 5.0 |
| Cabbage | 5.0 |
| Carrot | 7.0 |
| Cauliflower | 5.0 |
| Celeriac | 7.0 |
| Celery | 7.0 |
| Chard, Swiss | 7.5 |
| Chinese cabbage | 5.0 |
| Chives | 6.5 |
| Collards | 5.0 |
| Corn, sweet | 8.0 |
| Cucumber | 6.0 |
| Eggplant | 6.0 |
| Kale | 5.0 |
| Kohlrabi | 5.0 |
| Leek | 6.5 |
| Lettuce | 5.5 |
| Muskmelon | 6.0 |
| Mustard, India | 5.0 |
| Onion | 6.5 |
| Onion, Welsh | 6.5 |
| Parsley | 6.5 |
| Parsnip | 6.0 |

| Vegetable Seeds | Percent |
|---------------------------|---------|
| Pea | 7.0 |
| Pepper | 4.5 |
| Pumpkin | 6.0 |
| Radish | 5.0 |
| Rutabaga | 5.0 |
| Spinach | 8.0 |
| Squash | 6.0 |
| Tomato | 5.5 |
| Turnip | 5.0 |
| Watermelon | 6.5 |
| All other vegetable seeds | 6.0 |

(7-1-93)

04. Labeling. The container is conspicuously labeled in not less than 8-point type to indicate that the container is hermetically sealed, that the seed has been preconditioned as to moisture content, and the calendar month and year in which the germination test was completed. (7-1-93)

05. Germination. The percentage of germination of seed at the time of packaging was equal to or above the standards specified elsewhere in these rules. (7-1-93)

151. -- 199. (RESERVED).

200. NOXIOUS WEEDS.

01. Prohibited Noxious Weed Seeds -- Table 1.

| Common Name | Scientific Name |
|---------------------------|---|
| 1. Beancaper, Syrian | 1. <i>Zygophyllum fabago</i> L. |
| 2. Bindweed, Field | 2. <i>Convolvulus arvensis</i> L. |
| 3. Buffalobur | 3. <i>Solanum rostratum</i> Dun. |
| 4. Bursage, Skeleton leaf | 4. <i>Ambrosia tomentosa</i> Nutt. |
| 5. Camelthorn | 5. <i>Alhagi camelorum</i> Fisch. |
| 6. Carrot, Wild | 6. <i>Daucus carota</i> L. |
| 7. Cress, Hoary | 7. <i>Cardaria draba</i> (L.) Desv. |
| 8. Crupina, Common | 8. <i>Crupina vulgaris</i> Cass. |
| 9. Fieldcress, Austrian | 9. <i>Rorippa austriaca</i> (Crantz) Bess. |
| 10. Goatgrass | 10. <i>Aegilops cylindrica</i> Host |
| 11. Groundcherry, Smooth | 11. <i>Physalis subglabrata</i> Mackenz. & Bush |

| Common Name | Scientific Name |
|----------------------------|--|
| 12. Henbane, Black | 12. Hyoscyamus niger L. |
| 13. Johnsongrass | 13. Sorghum halepense L. |
| 14. Knapweed, Diffuse | 14. Centaurea diffusa Lam. |
| 15. Knapweed, Russian | 15. Centaurea repens L. |
| 16. Knapweed, Spotted | 16. Centaurea maculosa Lam. |
| 17. Lythrum, Purple | 17. Lythrum salicaria L. |
| 18. Nightshade, Silverleaf | 18. Solanum elaeagnifolium Cav. |
| 19. Pepperweed, Perennial | 19. Lepidium latifolium L. |
| 20. Poison-Hemlock | 20. Conium maculatum L. |
| 21. Puncturevine | 21. Tribulus terrestris L. |
| 22. Quackgrass | 22. Elytrigia repens; Agropyron repens (L.) Beauv. |
| 23. Ragwort, Tansy | 23. Senecio jacobaea L. |
| 24. Skeletonweed, Rush | 24. Chondrilla juncea L. |
| 25. Sowthistle, Perennial | 25. Sonchus arvensis L. |
| 26. Spurge, Leafy | 26. Euphorbia esula L. |
| 27. St. Johnswort, Common | 27. Hypericum perforatum L. |
| 28. Starthistle, Yellow | 28. Centaurea solstitialis L. |
| 29. Swainsonpea | 29. Sphaerophys salsula (Pall.) DC; Swainsona salsula (Pallas) Taubert |
| 30. Thistle, Canada | 30. Cirsium arvense (L.) Scop. |
| 31. Thistle, Musk | 31. Carduus nutans L. |
| 32. Thistle, Scotch | 32. Onopordum acanthium L. |
| 33. Toadflax, Dalmatian | 33. Linaria genistifolia spp. dalmatica (L.) Maire & Petit-mengin |
| 34. Toadflax, Yellow | 34. Linaria vulgaris Mill. |
| 35. Woad, Dyers | 35. Isatis tinctoria L. |

(7-1-93)

02. Restricted Noxious Weed Seeds -- Table 2.

| Common Name | Scientific Name |
|------------------|---|
| 1. Dodder | 1. Cuscuta spp. |
| 2. Halogeton | 2. Halogeton glomeratus (Stephen ex Bieb.) C.A. Mey |
| 3. Lettuce, Blue | 3. Lactuca pulchella (Pursch) DC. |

| Common Name | Scientific Name |
|-----------------------------------|---|
| 4. Oats, Wild | 4. Avena fatua L. |
| 5. Plantain, Buckhorn | 5. Plantago lanceolata L. |
| 6. Ragweed, Western and Perennial | 6. Ambrosia psilostachya DC., (T. & G.) Farw. |
| 7. Rye, Medusahead | 7. Taeniatherum caput-medusae (L.) Nevski; Elymus caput-medusae (L.) |
| 8. Sumpweed, Poverty | 8. Iva axillaris Pursh |

(7-1-93)

03. Restricted Noxious Weed Tolerances. (7-1-93)

a. Seven (7) seeds in fifty (50) grams of Agrotis species, Poa species, Rhodes grass, Bermudagrass, timothy, celery, and other agricultural or vegetable seeds of similar size and weight, within this group. (7-1-93)

b. Seven (7) seeds in each fifty (50) grams of Dallisgrass, ryegrass, fescue species, foxtail millets, alfalfa, red clover, sweetclover, lespedeza, bromegrass, Brassica species, carrot, onion, and other agricultural or vegetable seeds of similar size and weight or mixtures within this group, or mixtures of this group with those in group A. (7-1-93)

c. Seven (7) seeds in fifty (50) grams of alsike clover, white clover, and other agricultural or vegetable seeds of similar size and weight or mixtures within this group, with those in group A or group B. (7-1-93)

d. Eight (8) seeds in one hundred fifty (150) grams of Proso millet, Sudangrass, and seeds of similar size and weight, or mixtures of seed within this group. (7-1-93)

e. Forty-five (45) seeds in each pound for all wheatgrass species. (7-1-93)

f. Four (4) seeds in each five hundred (500) grams of wheat, oats, rye, barley, buckwheat, sorghums, vetches, field peas, and other seeds of a size and weight similar to or greater than those within this group, or any mixtures within this group. (7-1-93)

g. Containing more than one percent (1%) by weight of weed seeds including restricted noxious weed seeds referred to in Section 22-414(15), Idaho Code, provided, that three percent (3%) of cheat, chess, or downy brome shall be allowed in grass seed in which these weeds are found. (7-1-93)

201. -- 249. (RESERVED).

250. LABEL REQUIREMENTS OF SEEDS FOR SPROUTING.

The following information shall be indicated on all labels of seeds sold for sprouting in health food stores or other outlets: (7-1-93)

01. Name. Commonly accepted name of kind. (7-1-93)

02. Lot. Lot number. (7-1-93)

03. Percentage. Percentage by weight of the pure seed, crop seeds, inert matter, and weed seeds if required. (7-1-93)

04. Germination Percentage. Percentage of germination. (7-1-93)

05. Date. The calendar month and year the test was completed to determine such percentage. (7-1-93)

251. -- 299. (RESERVED).

300. VIABILITY BY TZ%.

A TZ (tetrazolium) test may be used in lieu of germination for the following species. The label shall read "viability by TZ%." (7-1-93)

01. Bitterbrush. (7-1-93)
02. Saltbush. (7-1-93)
03. Sagebrush. (7-1-93)
04. Indian ricegrass. (7-1-93)
05. Winterfat. (7-1-93)

301. -- 399. (RESERVED).

400. METHODS OF TESTING.

All methods used in testing and analyzing seed subject to this Act and the tolerances used in the enforcement of this Act shall conform as nearly as practicable to the current "Rules for Testing Seed adopted by the Association of Official Seed Analysts". (AOSA) file at the Idaho Department of Agriculture, State Seed Lab located at 2240 Kellogg Lane, Boise, Idaho 83712 and through the State Auditor's Office, Division of Statewide Rules, Office of the Rules Coordinator, located at 700 West State Street, Boise, Idaho 83720. (7-1-93)

401. -- 999. (RESERVED).